

Medicines and Doses to Reduce Pain and Fever in Children

Acetaminophen (Tylenol or other brand*): How much to give? Give every 4 to 6 hours, as needed. Do not give more than five doses in 24 hours.									
Child's Age	Infant's Liquid Syrup 160 mg/5 mL	Children's Liquid/Elixir Syrup 160 mg/5 mL	Children's Chewables 160 mg/tablet	Adult's Tablets 325 mg/tablet					
0 to 5 mos**	1.25 mL (40 mg)	1.25 mL (40 mg)	_	_					
6 to 11 mos	2.5 mL (80 mg)	2.5 mL (80 mg)	_	_					
12 to 23 mos	3.75 mL (120 mg)	3.75 mL (120 mg)	_	_					
2 to 3 years	5 mL (160 mg)	5 mL (160 mg)	1 tablet	_					
4 to 5 yrs	_	7.5 mL (240 mg)	1.5 tablets	_					
6 to 8 yrs	_	10 mL (320mg)	2 tablets	1 tablet					
9 to 10 yrs	_	12.5 mL (400 mg)	2.5 tablets	1 tablet					
11 yrs	_	15 mL (480 mg)	3 tablets	1.5 tablets					
12 yrs or older	_	_	4 tablets	2 tablets					
	O to 5 mos** 6 to 11 mos 12 to 23 mos 2 to 3 years 4 to 5 yrs 6 to 8 yrs 9 to 10 yrs 11 yrs	Eve every 4 to 6 hours, as needed. Description Child's Age Infant's Liquid Syrup 160 mg/5 mL 0 to 5 mos** 1.25 mL (40 mg) 6 to 11 mos 2.5 mL (80 mg) 12 to 23 mos 3.75 mL (120 mg) 2 to 3 years 5 mL (160 mg) 4 to 5 yrs — 9 to 10 yrs — 11 yrs —	Eve every 4 to 6 hours, as needed. Do not give more than fiven Child's Age Infant's Liquid Syrup 160 mg/5 mL Children's Liquid/Elixir Syrup 160 mg/5 mL 0 to 5 mos** 1.25 mL (40 mg) 1.25 mL (40 mg) 6 to 11 mos 2.5 mL (80 mg) 2.5 mL (80 mg) 12 to 23 mos 3.75 mL (120 mg) 3.75 mL (120 mg) 2 to 3 years 5 mL (160 mg) 5 mL (160 mg) 4 to 5 yrs - 7.5 mL (240 mg) 6 to 8 yrs - 10 mL (320mg) 9 to 10 yrs - 12.5 mL (400 mg) 11 yrs - 15 mL (480 mg)	Eve every 4 to 6 hours, as needed. Do not give more than five doses in 24 hours. Child's Age Infant's Liquid Syrup 160 mg/5 mL Children's Liquid/Elixir Syrup 160 mg/5 mL Children's Chewables 160 mg/tablet 0 to 5 mos** 1.25 mL (40 mg) - - 6 to 11 mos 2.5 mL (80 mg) - - 12 to 23 mos 3.75 mL (120 mg) - - 2 to 3 years 5 mL (160 mg) 5 mL (160 mg) 1 tablet 4 to 5 yrs - 7.5 mL (240 mg) 1.5 tablets 6 to 8 yrs - 10 mL (320mg) 2 tablets 9 to 10 yrs - 12.5 mL (400 mg) 2.5 tablets 11 yrs - 15 mL (480 mg) 3 tablets					

 $[^]st$ This table only includes products widely available in the United States.

^{**} Do not give acetaminophen to infants under two months old unless directed by healthcare provider.

Ibuprofen (Advil or Motrin*): How much to give? Give every 6 to 8 hours, as needed. Do not give more than four doses in 24 hours.									
Child's Weight	Child's Age	Infant's Drops/ Syringe 50 mg/1.25 mL	Children's Liquid/ Elixir 100 mg/5 mL	Children's Chewable Tablets 100 mg/tablet	Junior Strength Tablets 100 mg/tablet	Adult Strength Tablets 200 mg/tablet			
0 to 11 lbs 90 to 5 kg)	0 to 5 Months	DO NOT USE							
12 to 17 lbs (6 to 7 kg)	6 to 11 mos	1.25 mL (50 mg)	2.5 mL (50 mg)	_	_	_			
18 to 23 lbs (8 to 10 kg)	12 to 23 mos	1.875 mL (75 mg)	3.75 mL (75 mg)	_	_	_			
24 to 35 lbs (11 to 15 kg)	2 to 3 yrs	2.5 mL (100 mg)	5 mL (100 mg)	1 tablet (100 mg)	_	_			
36 to 47 lbs (16 to 21 kg)	4 to 5 yrs	_	7.5 mL (150 mg)	1.5 tablets (150 mg)	_	_			
48 to 59 lbs (22 to 26 kg)	6 to 8 yrs	_	10 mL (200mg)	2 tablets (200 mg)	2 tablets (200 mg)	1 tablet (200 mg)			
60 to 71 lbs (27 to 32 kg)	9 to 10 yrs	_	12.5 mL (250 mg)	2.5 tablets (250 mg)	2.5 tablets (250 mg)	1 tablet (200 mg)			
72 to 95 lbs (33 to 43 kg)	11 yrs	_	15 mL (300 mg)	3 tablets (300 mg)	3 tablets (300 mg)	1.5 tablets (300 mg)			
96 lbs (44 kg) or more	12 yrs or older	_	_	4 tablets (400 mg)	4 tablets (400 mg)	2 tablets (400 mg)			
* This table only includes products widely available in the United States.									

For dosage information or concerns, speak to an expert at Oklahoma Children's Hospital OU Health Pharmacy Phone: (405) 271-2156

